Draft version February 2, 2008
Preprint typeset using LaTeX style emulateapj v. 6/22/04

arXiv:astro-ph/0406595v1 25 Jun 2004

THE ANISOTROPY OF ELECTRON MHD TURBULENCE

Jungyeon Cho
CITA, Univ. of Toronto, 60 St. George St., Toronto, ON M5S 3H8, Canada; cho@cita.utoronto.ca
and
A. Lazarian
Astronomy Dept., Univ. of Wisconsin, Madison, WI53706, USA; lazarian@astro.wisc.edu
Draft version February 2, 2008

ABSTRACT

We present numerical studies of 3-dimensional electron magnetohydrodynamic (EMHD) turbulence. We investigate cascade timescale and anisotropy of freely decaying strong EMHD turbulence with zero electron skin depth. Cascade time scales with $k^{-4/3}$. Our numerical results clearly show scale-dependent anisotropy. We discuss that the observed anisotropy is consistent with $k_\parallel \propto k_\perp^{1/3}$, where $k_\parallel$ and $k_\perp$ are wave numbers parallel and perpendicular to (local) mean magnetic field, respectively.

Subject headings: MHD — turbulence — acceleration of particles

1. INTRODUCTION

Electron magnetohydrodynamics (EMHD) deals with MHD phenomena occurring at scales smaller than those of conventional MHD (see Kingsep, Chukbar, & Yankov 1990). On scales below the ion inertial length $d_i = c/\omega_{pi}$, where c is the speed of light and ω_{pi} is the ion plasma frequency, we can assume that the ions create only smooth motionless background and fast electron flows carry all the current, so that

$$\mathbf{v}_e = -\frac{\mathbf{J}}{ne} = -\frac{c}{4\pi ne}\nabla\times\mathbf{B}. \quad (1)$$

When we ignore the ambipolar diffusion, the Ohm's law becomes $\mathbf{E} = -\mathbf{v}_e\times\mathbf{B}/c + \eta\mathbf{J}$, where η is the plasma resistivity. Inserting this into Faraday's induction law ($\partial\mathbf{B}/\partial t = -c\nabla\times\mathbf{E}$), we obtain the EMHD equation

$$\frac{\partial\mathbf{B}}{\partial t} = -\frac{c}{4\pi ne}\nabla\times[(\nabla\times\mathbf{B})\times\mathbf{B}] + \eta\nabla^2\mathbf{B}. \quad (2)$$

Note that, in this paper, we only consider the zero (normalized) electron inertial length case: $d_e = c/(\omega_{pe}L) \rightarrow 0$, where ω_{pe} is the electron plasma frequency and L is the typical size of the system.

EMHD turbulence (also known as whistler turbulence) can be viewed as a generalized Alfvenic turbulence on scales smaller than the proton Larmor radius (Quataert & Gruzinov 1999). EMHD turbulence plays important roles in neutron stars and accretion disks. Goldreich & Reisenegger (1992) discussed the properties of EMHD turbulence and argued that EMHD turbulence can enhance ohmic dissipation of magnetic field in isolated neutron stars (see also Cumming, Arras, & Zweibel 2004). Quataert & Gruzinov (1999) discussed transition from conventional Alfvenic MHD cascade to EMHD cascade in advection dominated accretion flows (ADAF). Other applications of the EMHD model include quasi-collisionless magnetic reconnection in laboratory and space plasmas (Bulanov, Pegoraro, & Sakharov 1992; Biskamp, Schwarz, & Drake 1995; Avinash et al. 1998) and plasma opening switches and Z pinches (see Kingsep et al. 1990).

Earlier researchers convincingly showed that energy spectrum of EMHD turbulence is steeper than Kolmogorov's $k^{-5/3}$ spectrum. Using two-dimensional (2D) numerical simulations, Biskamp, Schwarz, & Drake (1996) found that energy spectrum follows $E(k) \propto k^{-\mu}$ with $\mu = 2.25 \pm 0.1$. They showed that the following Kolmogorov-type argument can explain this spectrum. Suppose that the eddy interaction time for eddies of size l is the usual eddy turnover time $t_{cas,l} \sim l/v_l$. Since $\mathbf{v} \propto \nabla\times\mathbf{B}$ (equation (1)), this becomes $t_{cas,l} \propto l^2/b_l$. Combining this with the constancy of spectral energy cascade rate ($b_l^2/t_{cas,l}$=constant), one obtains $E(k) \propto k^{-7/3}$. Note that $E(k)$ and b_l are related by $kE(k) \sim b_l^2$. Biskamp et al. (1999) also obtained a similar result for 3D. Ng et al. (2003) confirmed the 2D result of Biskamp et al. (1996). Therefore, earlier works strongly support that

$$E(k) \propto k^{-7/3} \quad (\text{as } d_e \rightarrow 0) \quad (3)$$

for both 2D and 3D. Recently, Galtier & Bhattacharjee (2003) studied energy spectrum of *weak* whistler turbulence.

Dastgeer et al. (2000), Ng et al. (2003), and Dastgeer & Zank (2003) numerically studied anisotropy of 2D EMHD turbulence. No 3D study has been reported yet. In this paper, we study anisotropy of 3D EMHD turbulence.

Method.— We have calculated the time evolution of decaying 3D incompressible electron magnetohydrodynamic turbulence. We have adopted a pseudospectral code to solve the normalized incompressible EMHD equation in a periodic box of size 2π:

$$\frac{\partial\mathbf{B}}{\partial t} = -\nabla\times[(\nabla\times\mathbf{B})\times\mathbf{B}] + \eta'\nabla^2\mathbf{B}, \quad (4)$$

where magnetic field, time, and length are normalized by a mean field B_0, the whistler time $t_w = L^2(\omega_{pe}/c)^2/\Omega_e$ (Ω_e= electron gyro frequency), and a characteristic length scale L (see, for example, Galtier & Bhattacharjee 2003). The resistivity η' in equation (4) is dimensionless. The dispersion relation of a whistler waves in this normalized units is $\omega = kk_\parallel B_0$, where $k_\parallel$ is the wave number parallel to the (local) mean magnetic field. The magnetic field consists of the uniform background field and a fluctuating field: $\mathbf{B} = \mathbf{B}_0 + \mathbf{b}$. The strength of the uniform background field, B_0, is set to 1. We use 288^3 collocation points. At $t = 0$, the random magnetic field



FIG. 1.— Energy spectra, timescale and anisotropy. (a) Energy spectra for t=0 (dotted), 0.33 (dashed), and 0.54 (solid). The spectra after $t > 0.33$ are compatible with $E(k) \propto k^{-7/3}$. (b) Cascade time scales with $t_{cas} \propto k^{-4/3}$. (c) Anisotropy calculated from equation (7).

is restricted to the range $2 \leq k \leq 4$ in wavevector ($\mathbf{k}$) space. The amplitudes of the random magnetic field at $t = 0$ is ~ 1. Hyperdiffusivity is used for the diffusion terms. The power of hyperdiffusivity is set to 3, so that the dissipation term in the above equation is replaced with $\eta_3(\nabla^2)^3\mathbf{B}$, where η_3 is approximately 2×10^{-10}.

2. RESULTS

Spectrum.—Figure 1 shows energy spectra. At $t = 0$ only large scale (i.e. small k) Fourier modes are excited. The dotted curve in Figure 1 shows the initial spectrum. As the turbulence decays, the initial energy cascades down to small scales and, as a result, small scale (i.e. large k) modes are excited. When the energy reaches the dissipation scale at $k > 70$, the energy spectrum decreases without changing its slope (the dashed and the solid curves). The slope at this stage is around 2.20, which is very close to the predicted spectrum:

$$E(k) \propto k^{-7/3}. \tag{5}$$

However, we also note that our result is compatible with a k^{-2} spectrum.

Timescale.—The energy spectra confirm that the Kolmogorov-type energy cascade model with $t_{cas} \sim l/v_l$ works fine for EMHD turbulence. Here we show that timescale of motion also supports the energy cascade model.

Symbolically, we can rewrite the EMHD equation in wave-vector space as follows: $\dot{\mathbf{b}}_{\mathbf{k}} = \mathbf{N}^b_{\mathbf{k}}$, where $\mathbf{N}^b_{\mathbf{k}}$ represents the nonlinear term. We have ignored the dissipation term. Naively speaking, we might obtain the time scale by dividing $|\mathbf{b}_{\mathbf{k}}|$ by $|\mathbf{N}^b_{\mathbf{k}}|$. However, to remove the effect of large scale translational motions, we restrict the evaluation of the nonlinear terms to contributions coming from the interactions between the mode at k and other modes within the range of $k/2$ and $2k$ (see Cho, Lazarian, & Vishniac (2002a) for details). We show the result in Figure 1(b). Our result seems to follow the timescale

$$t_{cas} \sim l/v_l \sim l^{4/3} \sim k^{-4/3}. \tag{6}$$

Anisotropy.— Since $\mathbf{B} \cdot \nabla\mathbf{B} \sim B_0 < k_{\|} > b$, we can obtain the average parallel wavenumber $< k_{\|} >$ from $< k_{\|} >\sim (|\mathbf{B} \cdot \nabla\mathbf{B}|^2/B_0^2/b^2)^{1/2}$, where b is the strength of the random magnetic field. The quantity $1/ < k_{\|} >$ is roughly the average parallel size of eddies at the energy injection scale. We can easily extend this method to smaller scale eddies. Consider eddies of perpendicular size l. Since it is reasonable to assume that the eddies are stretched along the *local* mean magnetic field, the quantity $\mathbf{B}_L \cdot \nabla\mathbf{b}_l$, not $\mathbf{B}_0 \cdot \nabla\mathbf{b}_l$, is proportional to $B_L k_{\|} b_l$, where $\mathbf{B}_L$ is the local mean field, $\mathbf{b}_l$ the fluctuating field at scale l, and $k_{\|}$ the wave number parallel to the local mean magnetic field (Cho, Lazarian, & Vishniac 2002b). Therefore, we have $k_{\|}(l) \sim (|\mathbf{B}_L \cdot \nabla\mathbf{b}_l|^2/B_L^2/b_l^2)^{1/2}$. It is convenient to do the calculation in Fourier space:

$$k_{\|}(k_{\perp}) \approx \left(\frac{\sum_{k \leq |\mathbf{k}'| < k+1} |\widehat{\mathbf{B}_L \cdot \nabla\mathbf{b}_l}|^2_{\mathbf{k}'}}{B_L^2 \sum_{k \leq |\mathbf{k}'| < k+1} |\hat{\mathbf{b}}|^2_{\mathbf{k}'}} \right)^{1/2}, \tag{7}$$

where $k_{\perp}$ ($\sim k \sim 1/l$ when anisotropy is present) is the wave number perpendicular to the local mean magnetic field and $\mathbf{B}_L$ is the local mean field obtained by eliminating modes whose perpendicular wavenumber is greater than $k/2$ ($\sim 1/(2l)$). The fluctuating field $\mathbf{b}_l$ is obtained by eliminating modes whose perpendicular wavenumber is less than $k/2$. We plot the result in Figure 1(c). The result does not show a well-defined power law between $k_{\|}$ and $k_{\perp}$. However, the result is roughly compatible with

$$k_{\|} \sim k_{\perp}^{1/3}. \tag{8}$$

We believe this is the true anisotropy of EMHD turbulence.

We may use the second order structure function to illustrate scale-dependent anisotropy of eddy structures (see Cho & Vishniac 2000). We calculate the second order structure function in a local frame, which is aligned with local mean magnetic field $\mathbf{B_L}$:

$$\mathrm{SF}_2(r_{\|}, r_{\perp}) =< |\mathbf{B}(\mathbf{x}+\mathbf{r}) - \mathbf{B}(\mathbf{x})|^2 >_{avg.\ over\ \mathbf{x}}, \tag{9}$$

where $\mathbf{r} = r_{\|}\hat{\mathbf{r}}_{\|} + r_{\perp}\hat{\mathbf{r}}_{\perp}$ and $\hat{\mathbf{r}}_{\|}$ and $\hat{\mathbf{r}}_{\perp}$ are unit vectors parallel and perpendicular to the local mean field $\mathbf{B_L}$, respectively. See Cho et al. (2002a) and Cho & Vishniac (2000) for the detailed discussion of the local frame. We show the resulting structure function in Figure 2.

We may obtain anisotropy of eddy structures by analyzing the contour shapes. However, we believe that this method will not give the true anisotropy for EMHD. The reason is as follows. Roughly speaking, when energy spectrum is steeper than k^{-3} we should not use the method. When energy spectrum is steeper than k^{-3},



FIG. 2.— Contour plot of the second structure function $SF_2(r_{\|}, r_\perp)$ at $t = 0.44$, where $r_{\|}$ ($r_\perp$) is distance along (perpendicular to) the *local* mean field, shows existence of scale-dependent anisotropy: smaller eddies are more elongated. Although this plot is useful to illustrate scale-dependent anisotropy, we do not use this plot to study anisotropy (see text).


B
$l_{||}$
$l_\perp$
$\mathbf{B}_L$
$\mathbf{b}_l$
$\frac{l_\perp}{l_{||}} = \frac{b_l}{B_L} \sim \frac{b_l}{B_0}$

FIG. 3.— Typical eddy shape and b_l/B_L ratio. The degree of bending of magnetic field lines depends on the b_l/B_L ratio. When we assume that $l_\perp$ and $l_{\|}$ represent typical perpendicular and parallel size of eddies at the scale l ($\sim l_\perp$), we obtain $l_\perp/l_{\|} \sim b_l/B_0$ and, thus, $l_{\|} \sim l_\perp^{1/3}$, we use $b_l \sim l^{2/3}$ and $B_L \sim B_0$.

small scale fluctuations are so small that the calculation of the second-order structure function will be dominated by the smooth change of the large scale field and the second-order structure function will show r^2-scaling regardless of the true scaling: $< |B(x) - B(x+r)|^2 > \sim < |B_L(x) - B_L(x+r)|^2 > \propto r^2$ when $r < L$. In the perpendicular direction, $E(k_\perp) \propto k_\perp^{-7/3}$ and, therefore, it is fine to use the structure function. However, we should not use the method for the *parallel* direction when anisotropy is stronger than $k_{\|} \propto k_\perp^{2/3}$. Suppose anisotropy scales with $k_{\|} \propto k_\perp^{\alpha}$. Then the 3D energy spectrum is $E_{3D}(k_{\|}, k_\perp) \propto k_\perp^{-10/3-\alpha} g(k_{\|}/k_\perp^{\alpha})$, where g is a function that describes distribution of energy along the $k_{\|}$ direction in Fourier space (see Goldreich & Sridhar 1995; Cho et al. 2002a). The 1D energy spectrum for the *parallel* direction becomes $E(k_{\|}) \propto \int k_\perp d\phi dk_\perp E_{3D}(k_{\|}, k_\perp) \propto k_{\|}^{-1-4/(3\alpha)}$ (Cho et al. 2002a). Therefore, when $\alpha < 2/3$, $E(k_{\|})$ is steeper than $k_{\|}^{-3}$ and, thus, we cannot use the structure function to study anisotropy. In other words, even if $\alpha < 2/3$, the structure function method will give a wrong result that $\alpha \approx 2/3$.

3. DISCUSSION

We obtained $k_{\|} \propto k_\perp^{1/3}$ in the previous section. See equation (8). Here we discuss why this is reasonable.

First, the fact that the Kolmogorov-type argument works fine for the energy spectrum implies the $k_{\|} \propto k_\perp^{1/3}$ anisotropy. As we mentioned earlier, Biskamp et al. (1996) assumed that the cascade time (t_{cas}) is the eddy turnover time ($t_{eddy} \sim l/v_l \sim l^2/b_l$) and obtained $E(k) \propto k^{-7/3}$, which agrees well with numerical simulations. See also our calculation of timescale in Figure 1(b). We can show that, when $t_{cas} \sim t_{eddy}$, there is balance between the eddy turnover time t_{eddy} and the the whistler time t_w (=the linear wave period=$\sim 1/(kk_{\|}B_0)$). Suppose that we have a wave packet whose parallel size is $l_{\|} \sim k_{\|}^{-1}$ and perpendicular size $l_\perp \sim k_\perp^{-1}$ ($\sim l \sim k^{-1}$ when anisotropy is present). This wave packet travels along the magnetic field line at the speed of kB_0. When this wave packet collides with opposite-traveling wave packets (of similar size), the change of energy per collision is $\Delta E \sim (dE/dt)\Delta t \sim (k^2 b_l^3)(l_{\|}/(kB_0))$, where we used equation (4) to estimate dE/dt. Therefore,

$$\Delta E/E \sim (kb_l)/(k_{\|}B_0) \sim t_w/t_{eddy}. \tag{10}$$

If $\Delta E/E < 1$, then many collisions are required to make $\Delta E/E \sim 1$. These collisions are incoherent. Then $(t_{eddy}/t_w)^2$ collisions are necessary to complete the cascade. This means that the energy cascade time by collision is

$$t_{cas,coll} \sim (t_{eddy}/t_w)^2 t_w. \tag{11}$$

If $t_{cas,coll} \sim t_{eddy}$, this equation implies $t_w \sim t_{eddy}$. Therefore, when $t_{cas} \sim t_{eddy}$, the turbulence is strong (i.e. $\Delta E/E \sim 1$) [1] and there is a balance between $t_{eddy} \sim 1/(k^2 b_l)$ and $t_w \sim 1/(kk_{\|}B_0)$. Our results clearly show that the critical balance factor $\xi \equiv t_w/t_{eddy}$ is constant (note that $b_l \sim l^{2/3}$ and $k_{\|} \propto k^{1/3}$). Using $b_l \sim k^{-2/3}$, we obtain $k_{\|} \propto k_\perp^{1/3}$. The uncertainty relation is another way to express this balance. The usual uncertainty relation between time and energy ($\sim$frequency) is $\Delta t \propto 1/\Delta\omega$. Since $\Delta t \sim t_{cas} \sim k^{-4/3}$ and $\Delta\omega \sim B_0 kk_{\|}$, we obtain the same anisotropy scaling.

Second, a simple geometric argument in Fig. 3 suggests the $k_{\|} \propto k_\perp^{1/3}$ anisotropy. Suppose that Fig. 3 shows a typical shape of magnetic field lines at scale l. Then, we can assume that $l_\perp/l_{\|} \sim b_l/B_L$, where b_l is the fluctuating field at the scale l and B_L is the local mean field ($B_L \sim B_0$). If we assume $l_\perp$ and $l_{\|}$ represent typical shape of eddies at the scale l ($\sim l_\perp$), we have

$$k_{\|}/k_\perp \sim b_l/B_0. \tag{12}$$

---

[1] The reverse is also true: if turbulence is strong, then $\Delta E/E \sim 1$ by definition and, therefore, energy cascade occurs during one wave period, which is equal to t_{eddy} (see equation (10)). Thus, one can show that three expressions, *Kolmogorov-type scaling for cascade time* ($t_{cas} \sim l/v_l$), *critical balance* ($t_{eddy} \sim t_w$), and *strong turbulence* ($\Delta E/E \sim 1$), are all equivalent (C. Thompson, private communication), as in *ordinary* MHD turbulence. See Goldreich & Sridhar (1995) for the original discussion of the critical balance and anisotropy of *ordinary* MHD turbulence.

In this interpretation, smaller eddies are more elongated because they have a smaller b_l/B_0 ratio. Note that the relation in equation (12) is in fact equivalent to the strong turbulence condition ($\Delta E/E \sim 1$; see equation (10)), which means energy cascade occurs over one wave period. From equation (10), we can show that $t_{cas} \sim l/v_l \sim l^2/b_l$. Combining this with the constancy of spectral energy cascade rate (b_l^2/t_{cas}=constant), we obtain $b_l \propto k^{-2/3}$ and, thus, the $k_\| \propto k_\perp^{1/3}$ anisotropy.

One peculiar feature of EMHD cascade is that there is balance between parallel and perpendicular cascade by self-interaction. The resonance relations of whistler turbulence,

$$\mathbf{k}_1 + \mathbf{k}_2 = \mathbf{k}_3 \quad \text{and} \quad k_1|k_{\|,1}| + k_2|k_{\|,2}| = k_3|k_{\|,3}|, \quad (13)$$

allow self-interaction. We can show that the parallel cascade by self interaction is as fast as the perpendicular cascade when $k_\| \propto k_\perp^{1/3}$. Ng et al. (2003) performed a numerical experiment and found that cascade by self-interaction is as fast as that by collision with opposite-traveling wave packets. Combined together, these facts imply that anisotropy cannot be stronger than $k_\| \propto k_\perp^{1/3}$ in EMHD. We will discuss this in detail in a forthcoming paper.

We have focused on anisotropy of 3D EMHD turbulence. However, it is fair to mention earlier 2D results. Dastgeer et al. (2000) numerically studied 2D EMHD turbulence and found numerical evidence that the turbulence is anisotropic in the presence of a mean field. Dastgeer & Zank (2003) showed that the *global* anisotropy of decaying EMHD turbulence scales linearly with the inverse of mean magnetic field strength, $1/B_0$. Ng et al. (2003) found local anisotropy of $k_\| \propto k_\perp^{3/4}$ from a 2D EMHD simulation. We will discuss the 2D versus 3D case elsewhere.

Conventional MHD turbulence can be decomposed into cascades of Alfven, fast and slow modes (Goldreich & Sridhar 1995; Cho & Lazarian 2002). While fast and slow modes get damped at larger scales, Alfvenic modes can cascade down to the proton Larmor radius scale. These Alfven modes excite whistler modes below the scale. Quataert & Gruzinov (1999) discussed transition from conventional Alfvenic MHD cascade to EMHD cascade in ADAFs and argued that anisotropy of EMHD turbulence is important for electron heating in ADAFs. They showed that, depending on the anisotropy of the resulting whistler turbulence, the whistler cascade will or will not transfer most of its energy to electrons. Whether or not whistler turbulence is anisotropic is an important question for the physics of accretion disks and gamma-ray bursts. The strong anisotropy of EMHD turbulence makes heating of protons by EMHD turbulence rather difficult in astrophysical plasmas with $\beta \sim 1$, where β is the gas to magnetic pressure ratio (see Quataert & Gruzinov 1999).

4. SUMMARY

Using a numerical simulation, we have studied 3D EMHD turbulence. We have found that energy cascade time scales with $k^{-4/3}$. We have calculated anisotropy and found $k_\| \propto k_\perp^{1/3}$. We have discussed why this is reasonable.

We thank Chris Thompson for many valuable comments and constructive suggestions. J.C. thanks Elliot Quataert for emphasizing the importance of the critical balance. This work was partially supported by NCSA under AST030023. A.L acknowledges NSF grant AST 0307869 ATM 0312282 and the NSF Center for Magnetic Self Organization in Laboratory and Astrophysical Plasmas.

REFERENCES

Avinash, K., Bulanov, S. V., Esirkepov, T., Kaw, P., Pegoraro, F., Sasorov, P. S., & Sen, A. 1998, Phys. Plasmas 5, 2849
Biskamp, D., Schwarz, E., & Drake, J. F. 1995, Phys. Rev. Lett. 75, 3850
Biskamp, D., Schwarz, E., & Drake, J. F. 1996, Phys. Rev. Lett., 76, 1264
Biskamp, D., Schwarz, E., Zeiler, A., Celani, A., & Drake, J. F. 1999, Phys. Plasmas, 6, 751
Bulanov, S. V., Pegoraro, F., & Sakharov, A. S. 1992, Phys. Fluids B, 4, 2499
Cho, J., Lazarian, A., & Vishniac, E. 2002a, ApJ, 564, 291
Cho, J., Lazarian, A., & Vishniac, E. 2002b, ApJ, 566, L49
Cho, J. & Lazarian, A. 2002, Phys. Rev. Lett., 88, 245001
Cho, J. & Vishniac, E. 2000, ApJ, 539, 273
Cumming, A., Arras, P., & Zweibel, E. 2004, ApJ, submitted (astro-ph/0402392)
Dastgeer, S., Das, A., Kaw, P., & Diamond, P. 2000, Phys. Plasmas, 7, 571
Dastgeer, S. & Zank, G. P. 2003, ApJ, 599, 715
Galtier, S. & Bhattacharjee, A. 2003, Phys. Plasmas, 10, 3065
Goldreich, P. & Reisenegger, A. 1992, ApJ, 395, 250
Goldreich, P. & Sridhar, S. 1995, ApJ, 438, 763
Kingsep, A. S., Chukbar, K. V., & Yan'kov, V. V. 1990, in *Reviews of Plasma Physics*, Vol. 16 (Consultants Bureau, New York)
Ng, C. S., Bhattacharjee, A., Germaschewski, K., & Galtier, S. 2003, Phys. Plasmas, 10, 1954
Quataert, E. & Gruzinov, A. 1999, ApJ, 520, 248